



# FRESH MARKET SWEET CORN

## Comparative table Sugary Enhanced, Synergistic

Disease resistance: R = Resistant - IR = Intermediate resistance

FRESH MARKET SWEET CORN	Type	Maturity (days)	Plant height	Ear height	Ear length	Ear diameter	Ear shape	Row count	Et	MDMV	Ps	Pst
<b>SUGARY ENHANCED</b>												
<b>Cloud Nine TSW®</b>	white se	76	80 in 203 cm	16 in 41 cm	8.5 in 21.5 cm			16-18	IR		IR	IR
<b>Colorow TSW®</b>	yellow se	77	74 in 188 cm	13 in 33 cm	8.5 in 21.5 cm			16-18	IR			IR
<b>Fastlane</b>	bicolor se	67	56 in 142 cm		7 in 18 cm			14-16				
<b>Lancelot</b>	bicolor se	82	80 in 203 cm	18 in 46 cm	9 in 23 cm			16-18	IR		IR	IR
<b>Luscious TSW®</b>	bicolor se	75	75 in 190 cm	15 in 38 cm	8 in 20.5 cm			16-18	IR		IR	IR
<b>Precious Gem</b>	bicolor se	80	75 in 190 cm	18 in 46 cm	9 in 23 cm			16-20	IR		IR	IR
<b>Spring Snow</b>	white se	67	42 in 107 cm	8 in 20.5 cm	7 in 18 cm			12-16	IR		IR	IR
<b>Spring Treat</b>	yellow se	68	60 in 152 cm	12 in 30.5 cm	8.5 in 21.5 cm			12	IR		IR	IR
<b>Sugar Pearl TSW®</b>	white se	73	78 in 198 cm	16 in 41 cm	8 in 20.5 cm			14-15	IR		IR	IR
<b>Tuxedo</b>	yellow se	78	72 in 183 cm	15 in 38 cm	8.5-9 in 21.5-23 cm			16-18	IR		IR	IR
<b>Welcome TSW®</b>	yellow se	72	72 in 183 cm	15 in 38 cm	8 in 20.5 cm			16-18	IR		IR	IR
<b>Whiteout</b>	white se	74	78 in 198 cm	16 in 41 cm	8 in 20.5 cm			16	IR			IR
<b>SYNERGISTIC</b>												
<b>Mattapoissett</b>	synergistic	80	78 in 198 cm	18 in 46 cm	8 in 20.5 cm			16-18	IR		IR	IR
<b>Misquamicut</b>	synergistic	78	78 in 198 cm	16 in 41 cm	8.5 in 21.5 cm			16-18	IR			IR
<b>Montauk</b>	synergistic	79	82 in 208 cm	18 in 46 cm	8 in 20.5 cm			16-18	IR			IR

Maturity: Relative maturities are based on 72% kernel moisture for su/se hybrids and 76% kernel moisture for sh2 hybrids. Maturity differences among the hybrids listed will remain fairly consistent. Actual maturities will vary depending on location and season.

*As with all sweet corn hybrids, certain herbicides may cause injury (phytotoxicity) to Bandero (HMX 6360) under certain conditions. Herbicide phytotoxicity has been observed during trialing of Bandero. Always consult the label for limitations and use caution at higher rates. Before applying any herbicide, take into account the stage of plant development, the soil type and condition, the temperature, moisture and wind. Injury may also result from the use of incompatible materials.*

Harris Moran Seed Company - PO Box 4938, Modesto, CA 95352, USA - Phone: 209.579.7333 - Web: [www.harrismoran.com](http://www.harrismoran.com)

*Data and disease ratings are an average of results from various test locations. It is not a prediction of future performance, but a summary of past results. Your performance will vary depending on the actual environment, pathogen strain and management conditions of your field. You must read and understand the Harris Moran Limitation of Warranty and Liability (found at [www.harrismoran.com](http://www.harrismoran.com)) before using this product.*

